

Product: **Animal Free Recombinant Human FGF-basic 147**
Cat #: 300-805P
Powder

Description	<p>Fibroblast Growth Factors (FGFs) are a 22 member family of proteins known to be involved in angiogenesis, wound healing and embryonic development. As a family, they bind to heparin and signal through four receptor tyrosine kinases called, FGFR1, 2, 3 and 4. Although the mechanism remains unclear, FGF-basic 147 (variant of FGF basic 154), also called FGF-2, is a critical component in keeping embryonic stem cells undifferentiated in cell culture systems. Alternate names: FGF2, HBGF-2, Prostatropin</p> <p>This product is produced with no animal-derived raw products, animal free equipment and animal free protocols.</p>
MW	Non-glycosylated protein, containing 147 amino acids, with a molecular weight of 16.5 kDa.
Physical Appearance	Sterile filtered white lyophilized (freeze-dried) powder.
Source	<i>E. coli</i>
Formulation	Recombinant human FGF-basic is lyophilized from a 10 mM Na ₂ PO ₄ , pH 8.0.
Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.
Stability	Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.
Biological Activity	The activity is determined by the dose-dependent proliferation of mouse BALB/c 3T3 cells and is typically less than 1 ng/mL.
Endotoxin Level	Measured by kinetic LAL analysis and is typically ≤ 1 EU/μg protein.
AA Sequence	MPALPEDGGS GAFPPGHFKD PKRLYCKNGG FFLRIHPDGR VDGVREKSDP HIKLQLQAE E RGVVSIKVC ANRYLAMKED GRLLASKCVT DECFFFERLE SNNYNTYRSR KYTSWYVALK RTGQYKLGSK TGPGQKAILF LPMSAKS

Purity greater than 97% determined by Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm.

Protein content determined by Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm.

THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!